

Teleportation scheme implementing the Universal Optimal Quantum Cloning Machine and the Universal NOT gate

M. Ricci, F. Sciarrino, C. Sias and F. De Martini

Dipartimento di Fisica and

Istituto Nazionale per la Fisica della Materia

Università di Roma "La Sapienza", Roma, 00185 - Italy

Abstract

The universal quantum cloning machine and the universal NOT gate acting on a single qubit can be implemented very generally by slightly modifying the protocol of quantum state teleportation. The experimental demonstration of the $1 \rightarrow 2$ cloning process according to the above scheme has been realized for a qubit encoded in photon polarization.

Classical information is encoded in bits, viz. dichotomic variables that can assume the values 0 or 1. For such variables there are no theoretical limitations as far as the "cloning" and/or "spin-flipping" processes are concerned. However manipulations on the quantum analogue of a bit, a qubit, have strong limitations due to fundamental requirements by quantum mechanics. For instance, it has been shown that an arbitrary unknown qubit cannot be perfectly cloned: $|\Psi\rangle \rightarrow |\Psi\rangle |\Psi\rangle$, a consequence of the so-called "no cloning theorem" [1], [2]. Another "impossible" device is the quantum NOT gate, the transformation that maps any qubit into the orthogonal one $|\Psi\rangle \rightarrow |\Psi^\perp\rangle$ [3]. In the last years a great deal of theoretical investigation has been devoted to finding the best approximation allowed by quantum mechanics to these two processes and to establish for them the corresponding "optimal" values of the "fidelity" $F < 1$. This problem has been solved in the general case [4–6]. In particular, it was found that a one-to-two universal optimal quantum cloning machine (UOQCM), i.e. able to clone one qubit into two qubits ($1 \rightarrow 2$), can be realized with

a fidelity $F_{CLON} = \frac{5}{6}$. It has also been proposed to test this result by a quantum optical "amplification" method, i.e. by associating the cloning effect with the QED "stimulated emission" process [7,8]. This proposal was quickly followed by some successful experimental demonstrations [9–11]. Very recently it has been argued by [11] that when the cloning process is realized in a subspace H of a larger nonseparable Hilbert space $H \otimes K$ which is acted upon by a physical apparatus, the same apparatus performs contextually in the space K the "flipping" of the input injected qubit, then realizing a $(1 \rightarrow 1)$ Universal-NOT gate (U-NOT) with a fidelity $F_{NOT} = \frac{2}{3}$ [5,6]. As an example, a UOQCM can be realized on one output mode of a non-degenerate "quantum-injected" optical parametric amplifier (QIOPA), while the U-NOT transformation is realized on the other mode [12]. Precisely this QIOPA apparatus has been adopted very recently to demonstrate experimentally the simultaneous, contextual realization of both processes [11,12].

In the present work this relevant, somewhat intriguing result is investigated under a new perspective implying a modified quantum state teleportation (QST) protocol, according to the following scheme. The QST protocol implies that an unknown input qubit $|\phi\rangle_S = \alpha|0\rangle_S + \beta|1\rangle_S$ is destroyed at a sending place (Alice: A) while its perfect replica appears at a remote place (Bob: B) via dual quantum and classical channels. To accomplish that two qualitatively different resources are needed: a quantum resource that cannot be cloned, and a classical resource that cannot travel faster than light [13,14]. Let us assume that Alice and Bob share the entangled "singlet" state $|\Psi^-\rangle_{AB} = 2^{-\frac{1}{2}}(|0\rangle_A|1\rangle_B - |1\rangle_A|0\rangle_B)$, and that we want to teleport the generic qubit $|\phi\rangle_S$. The "singlet" state is adopted here and hereafter because its well known invariance under $SU(2)$ transformations will ensure the "universality" of the cloning and U-NOT processes, as we shall see [5–8,11,12]. The overall state of the system is then $|\Omega\rangle_{SAB} = |\phi\rangle_S |\Psi^-\rangle_{AB}$. Alice performs a Bell measurement by projecting the qubits S and A into the four Bell states $\{|\Psi^-\rangle_{SA}, |\Psi^+\rangle_{SA}, |\Phi^-\rangle_{SA}, |\Phi^+\rangle_{SA}\}$ and then sends the result to Bob by means of 2 bits of classical information. In order to obtain $|\phi\rangle_B$, Bob applies to the received state the appropriate unitary transformation U_B according to the following protocol: $|\Psi^-\rangle_{SA} \rightarrow U_B = \mathbb{I}$, $|\Psi^+\rangle_{SA} \rightarrow U_B = \sigma_Z$, $|\Phi^-\rangle_{SA} \rightarrow U_B = \sigma_X$,

$|\Phi^+\rangle_{SA} \rightarrow U_B = \sigma_Y$ where the kets express the received corresponding information, I, σ_Z, σ_X are respectively the identity, phase flip, spin flip operators and $\sigma_Y = -i\sigma_Z\sigma_X$. The teleportation channel, after the unitary operation U_B , acts as the identity operator: $E_{QST}(\rho_S) = \rho_B$ where ρ_S is the density matrix representing the state of the qubit S . In absence of the classical information channel, Bob cannot apply the appropriate U_B , and the apparatus realized the map $E_B(\rho_S) = \frac{\mathbb{I}_B}{2}$, corresponding to the depolarizing channel E_{DEP} . This is the worst possible information transfer because any information about the initial state $|\phi\rangle$ is lost.

In order to implement the UOQCM and UNOT machines at Alice's and Bob's sites respectively, in the present letter we modify the QST protocol by performing a different projective measurement on the systems S and A . This leads to a different classical information to be transferred from A to B : Fig.1. Precisely, the complete measurement able to discriminate between the four Bell states is replaced by a dichotomic measurement able to identify $|\Psi^-\rangle_{SA}$ and its complementary space. Let us analyze the outcomes of such strategy. With a probability $p = \frac{1}{4}$ the state $|\Psi^-\rangle_{SA}$ is detected by A . In this case the correct QST channel E_{QST} is realized. If with probability $p = \frac{3}{4}$ this is not the case, Bob does not apply any unitary transformation to the set of non identified Bell states $\{|\Psi^+\rangle_{SA}, |\Phi^-\rangle_{SA}, |\Phi^+\rangle_{SA}\}$, and then the QST channel implements the statistical map $E(\rho) = \frac{1}{3}[\sigma_Z\rho\sigma_Z + \sigma_X\rho\sigma_X + \sigma_Y\rho\sigma_Y]$. This map indeed coincides with the map $E_{UNOT}(\rho_S)$ which realizes the Universal Optimal NOT gate, i.e. the one that approximates "optimally" the flipping of one qubit $|\phi\rangle$ into the orthogonal qubit $|\phi^\perp\rangle$. Bob identifies the two different maps realized at his site by reading the information (1 bit) received by Alice on the classical channel. For example, such bit can assume the value 0 if Alice identifies the Bell state $|\Psi^-\rangle_{SA}$ and 1 if not.

As already stated, in a bipartite entangled system the U-NOT gate is generally realized with an optimal quantum cloning process [11,12]. Therefore it is worth analyzing carefully what happens when the overall state $|\Omega\rangle_{SAB}$ is projected onto the subspace orthogonal to $|\Psi^-\rangle_{SA} \langle\Psi^-|_{SA} \otimes H_B$, i.e. by the projector:

$$P_{SAB} = (\mathbb{I}_{SA} - |\Psi^-\rangle_{SA} \langle\Psi^-|_{SA}) \otimes \mathbb{I}_B \quad (1)$$

It generates the normalized projected state $|\tilde{\Omega}\rangle \equiv P_{SAB}|\Omega\rangle_{SAB} = \sqrt{\frac{2}{3}}[|\xi_1\rangle_{SA} \otimes |1\rangle_B - |\xi_0\rangle_{SA} \otimes |0\rangle_B]$ where $|\xi_1\rangle_{SA} = \alpha|0\rangle_S|0\rangle_A + \frac{\beta}{2}|1\rangle_S|0\rangle_A + \frac{\beta}{2}|0\rangle_S|1\rangle_A$ and $|\xi_0\rangle_{SA} = \beta|1\rangle_S|1\rangle_A + \frac{\alpha}{2}|1\rangle_S|0\rangle_A + \frac{\alpha}{2}|0\rangle_S|1\rangle_A$. By projecting this state on the manifold SA the following density matrix is obtained:

$$\rho_{SA} \equiv Tr_B |\tilde{\Omega}\rangle \langle \tilde{\Omega}| = \frac{2}{3} |\phi\rangle \langle \phi|_{SA} + \frac{1}{3} |\{\phi, \phi^\perp\}\rangle_{SA} \langle \{\phi, \phi^\perp\}|_{SA} \quad (2)$$

where $|\{\phi, \phi^\perp\}\rangle_{SA} = 2^{-\frac{1}{2}}(|\phi^\perp\rangle_S|\phi\rangle_A + |\phi\rangle_S|\phi^\perp\rangle_A)$. We further project on the reduced spaces obtaining the matrices:

$$\rho_S \equiv Tr_A \rho_{SA} = \frac{5}{6} |\phi\rangle \langle \phi| + \frac{1}{6} |\phi^\perp\rangle \langle \phi^\perp| = \rho_A \equiv Tr_S \rho_{SA} \quad (3)$$

$$\rho_B \equiv Tr_{SA} |\tilde{\Omega}\rangle \langle \tilde{\Omega}| = \frac{2}{3} |\phi^\perp\rangle \langle \phi^\perp| + \frac{1}{3} |\phi\rangle \langle \phi| \quad (4)$$

by which the "optimal" values for the fidelities of the two "forbidden" processes are obtained: $F_{CLON} = \frac{5}{6}$ and $F_{UNOT} = \frac{2}{3}$ [11,12].

We want also to emphasize that it is possible to obtain the same results by performing a projective measurement on different Bell states. It's easily verified that by projection along $(\mathbb{I}_{SA} - |\Phi^+\rangle_{SA} \langle \Phi^+|_{SA}) \otimes \mathbb{I}_B$, the teleportation channel realizes the optimal universal approximation of the transpose map. This map is related to the UNOT map by a unitary transformation: $E_{TRANSPOSE} = E_{\sigma_Y}(E_{UNOT}(\rho))$, where $E_{\sigma_Y}(\rho) = \sigma_Y \rho \sigma_Y$. It's straightforward verified that the "cloning machine", in this case, generates $\rho'_S = \rho'_A = \sigma_Y \rho_S \sigma_Y$, that is, the clones of the input qubit on which acts the unitary transformation σ_Y . Similarly, by choosing another Bell state in (1), analogue transformations are achieved. Of course, in this case the universality of the processes is retrieved by applying an adequate unitary transformation to the output states of qubits labelled by SA and B .

Note that in the present context the presence of the entangled state $|\Psi^-\rangle_{AB}$ is not strictly necessary for the implementation of the solely quantum cloning process. Indeed for

this purpose we could inject into the Alice apparatus in Fig.2 the input qubit $|\phi\rangle$ together with a *fully mixed* state $\rho_A = \frac{\mathbb{I}_A}{2}$ spanning a 2 dimension space. Of course in this case only the Alice's apparatus is relevant and the U-Not process is absent. This is the solution chosen for our experimental demonstration: Fig.2. The qubit to be cloned is: $|\phi\rangle_S = \alpha|H\rangle_S + \beta|V\rangle_S$ where $|H\rangle$ and $|V\rangle$ respectively correspond to the horizontal and vertical linear polarizations of a single photon injected in one input mode S of a 50 : 50 beamsplitter BS_A . A fully mixed state ρ_A is simultaneously injected on the other input mode A of BS_A where the two input modes are linearly superimposed. Consider the overall output state which is realized on the two output modes 1 and 2 of BS_A . It can be expressed as a linear combination of the Bell states $\{|\Psi^-\rangle_{SA}, |\Psi^+\rangle_{SA}, |\Phi^-\rangle_{SA}, |\Phi^+\rangle_{SA}\}$. As it is well known, the realization of the singlet $|\Psi^-\rangle_{SA}$ is unambiguously identified by the detection of two single photons on the output modes of BS_A while the realization of the set of the other three Bell states correspond to the emission of photon pairs on either one of the output modes [15]. Hence the detection of two photons over either the mode 1 or 2, a Bose-mode-occupation enhancement, implies the projection by $P_{SA} = (\mathbb{I}_{SA} - |\Psi^-\rangle_{SA}\langle\Psi^-|)$ of the output state into the space orthogonal to $|\Psi^-\rangle_{SA}$ [16]. As seen in the theoretical analysis above this condition leads, in the general case, to the simultaneous realization of UOQCM and UNOT by a post-selection technique.

In the present experiment, a pair of non entangled photons with wavelength $\lambda = 532nm$ and with a coherence-time $\tau_{coh} = 80fs$, were generated by a spontaneous parametric down conversion SPDC process in a Type I BBO crystal in the initial polarization product state $|H\rangle_S |H\rangle_A$. The two photons were then injected on the two input modes S and A of BS_A . The input qubit $|\phi\rangle_S$, associated with mode S was polarization encoded by means of a waveplate (wp) WP_S . The transformation used to map the state $|H\rangle_A$ into $\rho_A = \frac{\mathbb{I}_A}{2}$ was achieved by stochastically rotating a $\lambda/2$ waveplate inserted on the mode A during the experiment. In this way the statistical evolution of $|H\rangle_A$ into two orthogonal states with equal probability was achieved. The photons S and A were injected in the two input arms of BS_A with a mutual delay Δt micrometrically adjustable by a translation stage with position settings $Z = 2\Delta tc$. The setting value $Z = 0$ was assumed to correspond to the full

overlapping of the photon pulses injected into BS_A , i.e. to the maximum photon interference.

For the sake of simplicity, we only analyzed the measurements performed on the BS_A output mode 1: Fig.2. The polarization state on this mode was analyzed by the combination of the wp WP_C and of the polarization beam splitter PBS_C . For each input polarization state $|\phi\rangle_S$, WP_C was set in order to make PBS_C to transmit $|\phi\rangle_S$ and reflect $|\phi^\perp\rangle_S$. The "cloned" state $|\phi\phi\rangle_S$ could be detected on mode 1 by a two-photon counter, realized in our case by first separating the two photons by a 50 : 50 beam splitter BS_C and then detecting the coincidence $[D_{A1}, D_{A2}]$ between the output detectors D_{A1} and D_{A2} : Fig.2. Any coincidence between detectors D_B and D_{A2} corresponded to the realization of the state $|\phi\phi^\perp\rangle_S$. First consider the cloning machine switched off, by setting: $\Delta t > \tau_{coh}$ i.e. by making S and A not interfering on BS_A . In this case, since the states $|\phi\phi\rangle_S$ and $|\phi\phi^\perp\rangle_S$ were realized with the same probability on mode 1, the rate of coincidences detected by $[D_{A1}, D_{A2}]$ and $[D_{A2}, D_B]$ were expected to be equal. By turning on the cloning machine, i.e. by setting $\Delta t \ll \tau_{coh}$, on mode 1 the output density matrix ρ_{SA} was realized implying an enhancement by a factor $R = 2$ of the counting rate by $[D_{A1}, D_{A2}]$ and no rate enhancement by $[D_{A2}, D_B]$. The measurement of R was carried out by coincidence measurements involving simultaneously $[D_{A1}, D_{A2}]$ and $[D_{A2}, D_B]$. The experimental data are reported in Figure 3 for three different input states $|\phi\rangle_S = |H\rangle$, $2^{-\frac{1}{2}}(|H\rangle + |V\rangle)$, $2^{-\frac{1}{2}}(|H\rangle + i|V\rangle)$. There circles and square marks refer respectively to the $[D_{A1}, D_{A2}]$ and $[D_{A2}, D_B]$ coincidences versus the position setting Z . We may check that the cloning process only affects the $|\phi\phi\rangle_S$ component, as expected and R is determined as the ratio between the peak values (cloning machine switched on) and the basis values (cloning machine off). The corresponding experimental values of the *cloning fidelity* $F = \frac{2R+1}{2R+2}$ are: $F_H = 0.827 \pm 0.002$; $F_{H+V} = 0.825 \pm 0.002$; $F_{H+iV} = 0.826 \pm 0.002$. These ones are to be compared with the optimal value $F_{th} = 5/6 \approx 0.833$ which corresponds to the limit value of the amplification ratio $R = 2$. As a lucky aspect of the present work, note the relevance in the theory above of the "singlet" state $|\Psi_{SA}^-\rangle$ in our measurement procedure. As already remarked, according to quantum theory the exclusive SU(2) invariance of this state should imply the insensitivity of the value of the

"fidelity" F to the photon polarization, i.e. to the input state $|\phi\rangle_S$. Indeed the experimental results just reported expressing very close values of F for different polarization states confirm the "*universality*" of the cloning process.

In summary, two relevant quantum information processes, forbidden by quantum mechanics in their exact form are found to be connected by a modified quantum state teleportation scheme and can be "optimally" realized by a simple linear method. As a partial implementation of the new protocol, an experimental demonstration of one to two universal quantum cloning has been performed by a quantum optical method. The results are found in good agreement with the theoretical expectations. We believe that the new perspective established by the present work will substantially enlighten an important and still unclear aspect of fundamental quantum theory.

This work has been supported by the FET European Network on Quantum Information and Communication (Contract IST-2000-29681: ATESIT), by Istituto Nazionale per la Fisica della Materia (PRA "CLON") and by Ministero dell'Istruzione, dell'Università e della Ricerca (COFIN 2002).

REFERENCES

- [1] D. Dieks, *Phys. Lett. A* **92**, 271 (1982)
- [2] W.K. Wootters, and W.H. Zurek, *Nature (London)* **299**, 802 (1982). The first no-cloning argument was expressed earlier by G.Ghirardi in a Referee Report for Foundations of Physics (1981) (private comm.).
- [3] H. Bechmann-Pasquinucci and N. Gisin, *Phys. Rev. A* **59**, 4238 (1999).
- [4] V. Bužek, and M. Hillery, *Phys. Rev. A* **54**, 1844 (1996).
- [5] R.F. Werner, *Phys. Rev. A* **58**, 1827 (1998); V. Bužek, M. Hillery, and R.F. Werner, *Phys. Rev. A* **60**, R2626 (1999). R. Derka, V. Buzek and A. Ekert, *Phys. Rev. Lett.* **80**, 1571 (1998).
- [6] N. Gisin, and S. Popescu, *Phys. Rev. Lett.* **83**, 432 (1999).
- [7] F. De Martini, V. Mussi, and F. Bovino, *Opt. Comm.* **179**, 581 (2000).
- [8] C. Simon, G. Weihs, and A. Zeilinger, *Phys. Rev. Lett.* **84**, 2993 (2000).
- [9] A. Lamas-Linares, C. Simon, J.C. Howell, and D. Bouwmeester, *Science* **296**, 712 (2002).
- [10] S. Fasel, N. Gisin, G. Ribordy, V. Scarani, and H. Zbinden, *Phys. Rev. Lett.* **89**, 107901 (2002)
- [11] D. Pelliccia, V. Schettini, F. Sciarrino, C. Sias, and F. De Martini, quant-ph/0302087
- [12] F. De Martini, V. Bužek, F. Sciarrino, and C. Sias, *Nature (London)* **419**, 815 (2002).
- [13] C. Bennett, G. Brassard, C. Crepeau, R. Jozsa, A. Peres, and W. Wootters, *Phys. Rev. Lett.* **70**, 1895 (1993).
- [14] C.H. Bennett, and D. Di Vincenzo, *Nature (London)* **404**, 247 (2000).
- [15] D. Bouwmeester, A. Ekert, A. Zeilinger, *The Physics of Quantum Information* (Springer,


2000).


[16] J.C. Howell, I.A. Khan, D. Bouwmeester, and N.P. Bigelow, quant-ph/0301170

Figure.1. General scheme for the simultaneous realization of the Universal Quantum Cloning Machine (UOQCM) and of the UNOT gate.




Figure.2. Experimental setup for the optical implementation of the UOQCM by a modified Teleportation protocol (left part of the Figure).

Figure.3. Experimental result of the universal cloning process for different input qubits corresponding to the encoded polarizations: $|H\rangle$, $2^{-1/2}(|H\rangle + |V\rangle)$ and $2^{-1/2}(|H\rangle + i|V\rangle)$.





Coincidence rate $\times 10^3 \text{ (s}^{-1}\text{)}$



Beamsplitter displacement (μm)